

Condition of Oak in Ornamental Plantings in Tashkent

N. Dubenok

Russian State Agrarian University – Moscow Agricultural Academy named after K.A. Timiryazeva, Doctor of Agricultural Sciences, Academician RAS, Head of Department Moscow, Russian Federation

Yu. Gninenko

Russian State Agrarian University – Moscow Agricultural Academy named after K.A. Timiryazev; Russian Research Institute of Forestry and Forestry Mechanization, Candidate of Biological Sciences, Head of Laboratory, Pushkino, Moscow Region, Russian Federation, gninenko-yuri@mail.ru

G. Larina

Russian Research Institute of Phytopathology, Doctor of Biol. Sci., professor, Head of the Laboratory, Bolshie Vyazemy, Moscow region, Russian Federation

O. Efimov

Russian State Agrarian University – Moscow Agricultural Academy named after K.A. Timiryazeva, Candidate of Agricultural Sciences, Associate Professor of the Department of Soil Science Moscow, Russian Federation

L. Seraya

Russian Scientific Research Institute of Phytopathology, Candidate of Biological Sciences, Head of the Department, Bolshie Vyazemy, Moscow region, Russian Federation

Key words: *English oak, soil conditions, harmful organisms*

A survey of the health of oak trees in urban plantations in the city of Tashkent was carried out, and it was found that damage to trees by some pathological factors is associated with changes in the state of the soil as a result of economic activities. English oak is an introduced species in the Central Asian region, and in recent years it has been exposed to dangerous phytophages in several places in the region. Urban management activities should not lead to worsening conditions for the growth of oak, as this weakens the trees and makes them more vulnerable to pests and diseases.

It is shown that the violation of soil and hydrological conditions, as a result of ill-conceived building, oppresses the oak and is the cause of the weakening of trees.